

SOBOLEV ORTHOGONAL POLYNOMIALS ON PRODUCT DOMAINS

LIDIA FERNÁNDEZ, FRANCISCO MARCELLÁN, TERESA E. PÉREZ,
MIGUEL A. PIÑAR, AND YUAN XU

ABSTRACT. Orthogonal polynomials on the product domain $[a_1, b_1] \times [a_2, b_2]$ with respect to the inner product

$$\langle f, g \rangle_S = \int_{a_1}^{b_1} \int_{a_2}^{b_2} \nabla f(x, y) \cdot \nabla g(x, y) w_1(x) w_2(y) dx dy + \lambda f(c_1, c_2) g(c_1, c_2)$$

are constructed, where w_i is a weight function on $[a_i, b_i]$ for $i = 1, 2$, $\lambda > 0$, and (c_1, c_2) is a fixed point. The main result shows how an orthogonal basis for such an inner product can be constructed for certain weight functions, in particular, for product Laguerre and product Gegenbauer weight functions, which serve as primary examples.

1. INTRODUCTION

Let $w_i(x)$ be a nonnegative weight function defined on an interval $[a_i, b_i]$, where $i = 1, 2$. Let W be the product weight function

$$(1.1) \quad W(x, y) := w_1(x)w_2(y), \quad (x, y) \in \Omega := [a_1, b_1] \times [a_2, b_2].$$

The purpose of this paper is to study orthogonal polynomials with respect to the inner product

$$(1.2) \quad \langle f, g \rangle_S = \iint_{\Omega} \nabla f(x, y) \cdot \nabla g(x, y) W(x, y) dx dy + \lambda f(c_1, c_2) g(c_1, c_2),$$

where $\lambda > 0$ and (c_1, c_2) is a fixed point, typically a corner point of the product domain Ω .

Sobolev orthogonal polynomials of one variable have been extensively studied (see the survey [7]). In particular, polynomials that are orthogonal with respect to the one-variable analogue of the inner product (1.1) were analyzed in [2]. In contrast, the study of such polynomials in several variables is a fairly recent affair. In [9], one of the earliest studies in several variables, Sobolev orthogonal polynomials with respect to an inner product similar to (1.2) on the unit ball of \mathbb{R}^d are constructed, where the discrete part could also be replaced by the integral on the

Date: May 15, 2014.

2000 Mathematics Subject Classification. 33C50, 42C10.

Key words and phrases. classical orthogonal polynomials, orthogonal polynomials in two variables, Sobolev inner products, product domain.

The work of the first, third and fourth author has been partially supported by DGICYT, Ministerio de Economía y Competitividad (MINECO) of Spain grant MTM 2011–28952–C02–02. The work of the second author has been supported by DGICYT, Ministerio de Economía y Competitividad (MINECO) of Spain grant MTM2012–36732–C03–01. The work of the fifth author was supported in part by NSF Grant DMS-1106113.

boundary of the ball. The motivation of [9] came from a question from engineering that requires control over the gradient. Such inner products appear naturally in the analysis of spectral methods for numerical solutions of partial differential equations (cf. [3]), which motivates our study.

For the ordinary inner product on the product domain,

$$(1.3) \quad \langle f, g \rangle_W = \iint_{\Omega} f(x, y)g(x, y) W(x, y) dx dy,$$

it is immediate that a basis of orthogonal polynomials of degree n is given by $p_k(w_1; x)p_{n-k}(w_2; y)$, $0 \leq k \leq n$, where $p_k(w; x)$ denotes the orthogonal polynomial of degree k with respect to w . A moment reflection shows, however, that Sobolev orthogonal polynomials with respect to the inner product (1.2) do not possess product structure. Our goal in this paper is to study the orthogonal structure for the inner product (1.2) on the product domain.

Our main result provides a way to construct a basis of Sobolev orthogonal polynomials, complemented with an algorithm that computes both orthogonal polynomials and their L^2 norm, when both weight functions w_1 and w_2 are self-coherent, which means that their monic orthogonal polynomials satisfy the relations of the form

$$(1.4) \quad p_n(x) = \frac{p'_{n+1}(x)}{n+1} + a_n p'_n(x) + b_n p'_{n-1}(x), \quad n \geq 1.$$

Weight functions, or measures, that are self-coherent have been studied extensively and characterized. They are essentially the classical measures. In [4] the authors proved that (1.4) characterizes classical orthogonal polynomials.

Our approach is to express the Sobolev orthogonal polynomials with respect to the inner product $\langle \cdot, \cdot \rangle_S$ in terms of a family of product polynomials, which are not, however, the product orthogonal polynomials with respect to (1.3), but product polynomials of the form $q_k(w_1; x)q_{n-k}(w_2; y)$, where $q_k(w)$ takes the form of the right hand side of (1.4) without the derivative. In order to keep the idea transparent, we will not work with the most general case that our method applies, but work primarily with two examples, product Laguerre weight functions and product Gegenbauer weight functions, for which we work out our algorithms explicitly.

Some of our results can no doubly be extended from two variables to several variables. We choose to stay with two variables to avoid complicated notation and keep the algorithm practical.

The paper is organized as follows. In the next section, we recall the basics for orthogonal polynomials of several variables, and describe our strategy for constructing Sobolev orthogonal polynomials for the product weight functions. The construction is worked out explicitly in the case of product Laguerre weight in Section 3 and in the case of product Gegenbauer weight in Section 4.

2. CONSTRUCTING BASES FOR SOBOLEV ORTHOGONAL POLYNOMIALS

The basics of orthogonal polynomials in several variables are given in the first subsection. Sobolev orthogonal polynomials for product measures are described in the second subsection, and the strategy for constructing an orthogonal basis is discussed in the third subsection.

2.1. Orthogonal polynomials of two variables. Let Π^2 denote the space of polynomials in two real variables and, for $n = 0, 1, 2, \dots$, let Π_n^2 denote the subspace of polynomials of (total) degree at most n in Π^2 . For an inner product $\langle \cdot, \cdot \rangle$ defined on Π^2 , a polynomial $P \in \Pi_n^2$ is said to be orthogonal if $\langle P, Q \rangle = 0$ for all $Q \in \Pi_{n-1}^2$. Let \mathcal{V}_n^2 denote the space of orthogonal polynomials of total degree n with respect to $\langle \cdot, \cdot \rangle$. It is known that

$$\dim \Pi_n^2 = \binom{n+2}{n} \quad \text{and} \quad \dim \mathcal{V}_n^2 = n+1.$$

The space \mathcal{V}_n^2 can have many different bases. A basis $\{P_k^n : 0 \leq k \leq n\}$ of \mathcal{V}_n^2 is called mutually orthogonal if $\langle P_k^n, P_j^n \rangle = 0$ for $k \neq j$ and it is called orthonormal if, in addition, $\langle P_k^n, P_k^n \rangle = 1$. Another polynomial basis that is of interest is the monic basis, for which $P_k^n(x, y) = x^{n-k}y^k + R_k^n(x, y)$, where $R_k^n \in \Pi_{n-1}^2$, $0 \leq k \leq n$. It is often convenient to use the vector notation

$$\mathbb{P}_n = (P_0^n, P_1^n, \dots, P_n^n)^\top,$$

considered as a column vector, which we also regard as a set of orthogonal polynomials of degree n . In this notation, $\langle \mathbb{P}_n, \mathbb{P}_m^\top \rangle = \mathbf{H}_n \delta_{n,m}$, where \mathbf{H}_n is a matrix of size $(n+1) \times (n+1)$, necessarily symmetric and positive definite. If the set \mathbb{P}_n contains a mutually orthogonal basis then \mathbf{H}_n is diagonal, and if it is orthonormal then \mathbf{H}_n is the identity matrix.

For $W(x, y) = w_1(x)w_2(y)$ as in (1.1), we consider the inner product

$$\langle f, g \rangle_W = c \int_{\Omega} f(x, y)g(x, y)W(x, y)dx dy,$$

where c is a normalization constant of W so that $\langle 1, 1 \rangle_W = 1$. A basis of $\mathcal{V}_n^2(W)$ is given by the product polynomials

$$(2.1) \quad P_k^n(x, y) := p_{n-k}(w_1; x)p_k(w_2; y), \quad 0 \leq k \leq n,$$

where $p_k(w_i; x) = x^k + \dots$ denotes the monic orthogonal polynomial with respect to w_i on $[a_i, b_i]$. Then P_k^n is the monic orthogonal polynomial and $\{P_k^n : 0 \leq k \leq n\}$ forms a mutually orthogonal basis of $\mathcal{V}_n^2(W)$.

2.2. Sobolev orthogonal polynomials. For $i = 1, 2$, let w_i be a weight function defined on the interval $[a_i, b_i]$, where $-a_i$ and b_i can be infinity. For the product weight function W in (1.1), let $\mathcal{V}_n^2(S)$ denote the space of Sobolev orthogonal polynomials of degree n with respect to the inner product $\langle \cdot, \cdot \rangle_S$ defined in (1.2). Most of our work will be carried out for the following two examples.

Example 2.1. For $\alpha > -1$, let w_α be the Laguerre weight function

$$w_\alpha(x) := x^\alpha e^{-x}, \quad x \in \mathbb{R}_+ := [0, \infty).$$

For $\alpha, \beta > -1$, let $W_{\alpha, \beta}$ be the product Laguerre weight function defined by

$$W_{\alpha, \beta}(x, y) := w_\alpha(x)w_\beta(y), \quad (x, y) \in \Omega := \mathbb{R}_+^2.$$

There is only one finite corner point of Ω , and we consider the inner product

$$(2.2) \quad \langle f, g \rangle_S = c_{\alpha, \beta} \int_{\mathbb{R}_+^2} \nabla f(x, y) \cdot \nabla g(x, y) W_{\alpha, \beta}(x, y) dx dy + \lambda f(0, 0)g(0, 0),$$

where $\lambda > 0$ is a fixed constant and $c_{\alpha, \beta} = 1 / \int_{\mathbb{R}_+^2} W_{\alpha, \beta}(x, y) dx dy$.

Example 2.2. For $\alpha > -1/2$, let u_α be the Gegenbauer weight function

$$u_\alpha(x) := (1 - x^2)^{\alpha - \frac{1}{2}}, \quad x \in [-1, 1].$$

For $\alpha, \beta > -1/2$, let $U_{\alpha, \beta}$ be the product Gegenbauer weight function defined by

$$U_{\alpha, \beta}(x, y) := u_\alpha(x)u_\beta(y), \quad (x, y) \in \Omega := [-1, 1]^2.$$

There are four corner points of Ω and we consider the inner product

$$(2.3) \quad \langle f, g \rangle_S = c_{\alpha, \beta} \int_{-1}^1 \int_{-1}^1 \nabla f(x, y) \cdot \nabla g(x, y) U_{\alpha, \beta}(x, y) dx dy + \lambda f(1, 1)g(1, 1),$$

where $\lambda > 0$ is a fixed constant and $c_{\alpha, \beta} = 1 / \int_\Omega U_{\alpha, \beta}(x, y) dx dy$.

For the inner product $\langle \cdot, \cdot \rangle_S$ in (1.2), we denote its main part by

$$(2.4) \quad \begin{aligned} \langle f, g \rangle_\nabla &:= c \int_\Omega \nabla f(x, y) \cdot \nabla g(x, y) W(x, y) dx dy \\ &= \langle \partial_1 f, \partial_1 g \rangle_W + \langle \partial_2 f, \partial_2 g \rangle_W. \end{aligned}$$

This is a bilinear form and it is an inner product on the linear space $\Pi^2 \setminus \mathbb{R}$ of polynomials having a zero constant term. Let

$$\mathcal{V}_n^2(S) := \mathcal{V}_n^2(S, W) \quad \text{and} \quad \mathcal{V}_n^2(\nabla) := \mathcal{V}_n^2(\nabla, W)$$

denote the linear spaces of orthogonal polynomials of total degree n associated with $\langle \cdot, \cdot \rangle_S$ and $\langle \cdot, \cdot \rangle_\nabla$, respectively.

Let S_k^n be the monic orthogonal polynomial of degree n in $\mathcal{V}_n^2(S)$ that satisfies $S_k^n(x, y) - x^{n-k}y^k \in \Pi_{n-1}^2$ for $0 \leq k \leq n$. Likewise, for $n \geq 1$, let S_k^n be a monic orthogonal polynomial in $\mathcal{V}_n^2(\nabla)$.

Theorem 2.3. For $n \geq 1$, let $\{S_k^n : 0 \leq k \leq n\}$ denote a monic orthogonal basis of $\mathcal{V}_n^2(\nabla)$. Then, the monic orthogonal basis $\{S_k^n : 0 \leq k \leq n\}$ of $\mathcal{V}_n^2(S)$ is given by $S_0^0(x, y) = 1$ and

$$S_k^n(x, y) = S_k^n(x, y) - S_k^n(c_1, c_2), \quad n \geq 1.$$

Proof. Since $S_k^n(c_1, c_2) = 0$, it follows that $\langle S_k^n, S_j^m \rangle_S = \langle S_k^n, S_j^m \rangle_\nabla$ if $n \geq 1$. \square

This theorem shows that we only need to work with the bilinear form $\langle \cdot, \cdot \rangle_\nabla$ and on the linear space $\Pi^2 \setminus \mathbb{R}$. Observe that the orthogonal polynomials in $\mathcal{V}_n^2(\nabla)$ are determined up to an additive constant c . Indeed, for any constant c , the polynomial $S_k^n + c$ is also a monic orthogonal polynomial in $\mathcal{V}_n^2(\nabla)$. By Theorem 2.3, however, we only need to determine S_k^n up to a constant. For convenience, we adopt the following notation for two functions that are equal up to a constant:

$$f(x, y) \stackrel{c}{=} g(x, y) \quad \text{if} \quad f(x, y) - g(x, y) \equiv c,$$

where $c \in \mathbb{R}$ is a generic constant.

2.3. Strategy for constructing Sobolev orthogonal polynomials. In order to construct the polynomial S_k^n , we expand it in terms of a known basis of polynomials denoted by $\{Q_j^m : 0 \leq j \leq m \leq n\}$,

$$(2.5) \quad S_k^n(x, y) = \sum_{m=0}^n \sum_{j=0}^m a_{j,m}(k) Q_j^m(x, y),$$

and determine the coefficients $a_{j,m}(k)$ by orthogonality. Since S_k^n is determined up to a constant, the equal sign should be replaced by $\stackrel{c}{=}$ in (2.5).

The choice of Q_j^m clearly matters. An obvious choice is the basis of product orthogonal polynomials P_k^n in (2.1). This basis, however, is not a good choice since we need to work with derivatives of the basis elements. This is where the notion of coherent pair comes in.

A weight function w defined on the real line is called self-coherent if its monic orthogonal polynomials $p_n(w)$ satisfy the relation

$$(2.6) \quad p_n(w; x) = \frac{p'_{n+1}(w; x)}{n+1} + a_n p'_n(w; x), \quad n \geq 0,$$

for some constants a_n . Furthermore, w is called symmetric self-coherent, if w is an even function and its monic orthogonal polynomials $p_n(w)$ satisfy the relation

$$(2.7) \quad p_n(w; x) = \frac{p'_{n+1}(w; x)}{n+1} + b_n p'_{n-1}(w; x), \quad n \geq 1.$$

More generally, we can call w self-coherent if it satisfies (1.4), that is,

$$p_n(w; x) = \frac{p'_{n+1}(w; x)}{n+1} + a_n p'_n(w; x) + b_n p'_{n-1}(w; x), \quad n \geq 1$$

If w is self-coherent, we denote by $q_n(w)$ the polynomial of degree n defined by

$$(2.8) \quad q_n(w; x) = p_n(w; x) + n a_{n-1} p_{n-1}(w; x) + n b_{n-1} p_{n-2}(w; x), \quad n \geq 1,$$

where, by convention, $p_{-1}(w; x) = 0$ and we assume the last term is zero if $n = 1$. It follows directly from the definition that $q_n(w)$ is monic and

$$q'_n(w; x) = n p_{n-1}(w; x).$$

Notice that self-coherent orthogonal polynomials are essentially, up to a linear change of variable, the classical orthogonal polynomials (Jacobi, Laguerre and Hermite) as was proved in [4].

We now define the polynomials Q_j^m of two variables by

$$(2.9) \quad Q_k^n(x, y) := q_{n-k}(w_1; x) q_k(w_2; y), \quad 0 \leq k \leq n, \quad n = 0, 1, \dots$$

The derivatives of Q_k^n can be given explicitly in terms of product orthogonal polynomials P_j^m in (2.1).

Lemma 2.4. *Let ∂_i denote the i -th partial derivative. Then*

$$\begin{aligned} \partial_1 Q_0^n(x, y) &= n p_{n-1}(w_1; x) = n P_0^{n-1}(x, y) \quad \text{and} \quad \partial_2 Q_0^n(x, y) = 0, \\ \partial_1 Q_n^n(x, y) &= 0 \quad \text{and} \quad \partial_2 Q_0^n(x, y) = n p_{n-1}(w_2; y) = n P_{n-1}^{n-1}(x, y). \end{aligned}$$

Furthermore, for $1 \leq k \leq n-1$,

$$\begin{aligned} \partial_1 Q_k^n &= (n-k) \left(P_k^{n-1} + k a_{k-1}(w_2) P_{k-1}^{n-2} + k b_{k-1}(w_2) P_{k-2}^{n-3} \right), \\ \partial_2 Q_k^n &= k \left(P_{k-1}^{n-1} + (n-k) a_{n-k-1}(w_1) P_{k-1}^{n-2} + (n-k) b_{n-k-1}(w_1) P_{k-1}^{n-3} \right). \end{aligned}$$

Proof. For $1 \leq k \leq n$, it follows directly from the definition of Q_k^n that

$$\partial_1 Q_k^n(x, y) = q'_{n-k}(w_1; x) q_k(w_2; y) = (n-k) p_{n-k-1}(w_1; x) q_k(w_2; y).$$

Substituting $q_k(w_2; y)$ by its definition (2.8), the identity for $\partial_1 Q_k^n$ follows from the definition of P_j^m . The other identities are proved similarly. \square

Let $\mathbb{Q}_n = (Q_0^n, \dots, Q_n^n)^\top$ and $\mathbb{S}_n = (S_0^n, \dots, S_n^n)^\top$ denote the column vector of polynomials Q_k^n and S_k^n , respectively. Furthermore, let e_i denote the standard Euclidean coordinate vector whose i -th element is 1 and all other elements are 0.

Theorem 2.5. *For $0 \leq k \leq n$, there exist real numbers $a_{i,k}$ and $b_{i,k}$ such that*

$$(2.10) \quad Q_k^n(x, y) \stackrel{c}{=} S_k^n(x, y) + \sum_{i=0}^{n-1} a_{i,k} S_i^{n-1}(x, y) + \sum_{i=0}^n b_{i,k} S_i^{n-2}(x, y).$$

Moreover, in the case of $k = 0$ and $k = n$, we have, respectively,

$$(2.11) \quad S_0^n(x, y) \stackrel{c}{=} Q_0^n(x, y) \quad \text{and} \quad S_n^n(x, y) \stackrel{c}{=} Q_n^n(x, y).$$

In terms of vector notation, (2.10) can be written as

$$(2.12) \quad \mathbb{Q}_n \stackrel{c}{=} \mathbb{S}_n + \mathbf{A}_{n-1} \mathbb{S}_{n-1} + \mathbf{B}_{n-2} \mathbb{S}_{n-2},$$

where \mathbf{A}_{n-1} and \mathbf{B}_{n-2} are matrices of the form

$$\mathbf{A}_{n-1} = \begin{bmatrix} 0 & \dots & 0 \\ \hline & \tilde{\mathbf{A}}_{n-1} & \\ \hline 0 & \dots & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B}_{n-2} = \begin{bmatrix} 0 & \dots & 0 \\ \hline & \tilde{\mathbf{B}}_{n-2} & \\ \hline 0 & \dots & 0 \end{bmatrix}.$$

Here $\tilde{\mathbf{A}}_{n-1}$ and $\tilde{\mathbf{B}}_{n-2}$ are matrices of size $(n-1) \times n$ and $(n-1) \times (n-1)$, respectively.

Proof. If $k = 0$ and P is any polynomial in Π_{n-1}^2 , then, by Lemma 2.4,

$$\langle Q_0^n, P \rangle_{\nabla} = \langle P_0^{n-1}, \partial_1 P \rangle_W = 0.$$

Since the space $\{\partial_1 P : P \in \Pi_{n-1}^2\}$ is Π_{n-2}^2 , this shows that $Q_0^n \in \mathcal{V}_n^2(\nabla)$ and it is equal to S_0^n as it is monic. The proof for S_n^n is similar. Moreover, if $1 \leq k \leq n$, it follows from Lemma 2.4 that

$$\langle Q_k^n, P \rangle_{\nabla} = \langle \partial_1 Q_k^n, \partial_1 P \rangle_W + \langle \partial_2 Q_k^n, \partial_2 P \rangle_W = 0$$

for any polynomial P of degree at most $n-3$. Consequently, Q_k^n can be written as a linear combination of the Sobolev orthogonal polynomials of degree $n, n-1$ and $n-2$. Since both Q_k^n and S_k^n are monic by definition, (2.10) follows. \square

To determine the matrices \mathbf{A}_{n-1} and \mathbf{B}_{n-2} , we need to work with specific weight functions. The simplest cases are the product Laguerre polynomials for which $\mathbf{B}_{n-2} = 0$ and the product Gegenbauer polynomials for which $\mathbf{A}_{n-1} = 0$. These two cases will be worked out in detail in the next two sections.

3. THE PRODUCT LAGUERRE WEIGHT

In this section we consider the product of Laguerre weight functions and the inner product (2.2). The Laguerre polynomials are defined by (cf. [8, Chapt V])

$$L_n^\alpha(x) := \frac{(\alpha+1)_n}{n!} {}_1F_1(-n; \alpha+1; x) = \frac{(-1)^n}{n!} x^n + \dots$$

and their orthogonality is given by

$$\langle L_n^\alpha, L_m^\alpha \rangle_{w_\alpha} := \frac{1}{\Gamma(\alpha+1)} \int_0^{+\infty} L_n^\alpha(x) L_m^\alpha(x) w_\alpha(x) dx = \frac{(\alpha+1)_n}{n!} \delta_{n,m},$$

where $(a)_n = a(a+1)\cdots(a+n-1)$, $n \geq 1$, $(a)_0 = 1$, is the Pochhammer symbol. Furthermore, they satisfy the relation ([8, p. 102])

$$L_n^\alpha(x) = -\frac{d}{dx} L_{n+1}^\alpha(x) + \frac{d}{dx} L_n^\alpha(x),$$

which shows that the Laguerre weight function w_α is self-coherent. Monic Laguerre orthogonal polynomial $p_n(w_\alpha)$ and its L^2 norm are given by

$$p_n(w_\alpha; x) := (-1)^n n! L_n^\alpha(x), \quad h_n^\alpha := \langle p_n(w_\alpha), p_n(w_\alpha) \rangle_{w_\alpha} = n! (\alpha + 1)_n.$$

From these relations, it follows readily that the polynomial

$$q_n(w_\alpha; x) := p_n(w_\alpha; x) + np_{n-1}(w_\alpha; x)$$

satisfies $q_n'(w_\alpha; x) = np_{n-1}(w_\alpha; x)$ for $n = 0, 1, 2, \dots$

We are now ready to state our polynomials in two variables for the product Laguerre weight function $W_{\alpha, \beta}$ on \mathbb{R}_+^2 , with $\alpha, \beta > -1$. We again denote the orthogonal polynomials by P_k^n ,

$$P_k^n(x, y) := p_{n-k}(w_\alpha; x) p_k(w_\beta; y), \quad 0 \leq k \leq n.$$

It follows readily that these are mutually orthogonal polynomials and

$$(3.1) \quad h_k^n := \langle P_k^n, P_k^n \rangle_{W_{\alpha, \beta}} = h_{n-k}^\alpha h_k^\beta = (n-k)! k! (\alpha + 1)_{n-k} (\beta + 1)_k.$$

We also define the monic polynomial Q_k^n by

$$Q_k^n(x, y) := q_{n-k}(w_\alpha; x) q_k(w_\beta; y), \quad 0 \leq k \leq n.$$

In this setting, their partial derivative for $1 \leq k \leq n$ in Lemma 2.4 becomes the following:

Lemma 3.1. *For $1 \leq k \leq n-1$, the following formulas hold*

$$\begin{aligned} \partial_1 Q_k^n(x, y) &= (n-k) [P_k^{n-1}(x, y) + k P_{k-1}^{n-2}(x, y)], \\ \partial_2 Q_k^n(x, y) &= k [P_{k-1}^{n-1}(x, y) + (n-k) P_{k-1}^{n-2}(x, y)]. \end{aligned}$$

Recall that $\mathcal{V}_n^2(\nabla, W_{\alpha, \beta})$, $n \geq 1$, is the space of Sobolev orthogonal polynomials with respect to the bilinear form $\langle \cdot, \cdot \rangle_\nabla$ defined in (2.4). Let $S_k^n = x^{n-k} y^k + \dots$ be a monic orthogonal polynomial in $\mathcal{V}_n^2(\nabla, W_{\alpha, \beta})$. Then relation (2.12) becomes

$$(3.2) \quad \mathbb{Q}_n \stackrel{c}{=} \mathbb{S}_n + \mathbf{A}_{n-1} \mathbb{S}_{n-1}.$$

Our goal is to show how \mathbf{A}_{n-1} can be explicitly computed. To this end, we need explicit formulas for the inner products of the gradients of the polynomials Q_k^n . In the following we write $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_{W_{\alpha, \beta}}$.

Lemma 3.2. *For $0 \leq i \leq n$ and $0 \leq l \leq m$,*

$$\begin{aligned} \langle Q_i^n, Q_l^m \rangle_\nabla &= [l(m-l)^2 h_{l-1}^{m-2} \delta_{i, l-1} + l^2(m-l) h_{l-1}^{m-2} \delta_{i, l}] \delta_{n, m-1} \\ &\quad + [(m-l)^2 h_l^{m-1} \delta_{i, l} + 2l^2(m-l)^2 h_{l-1}^{m-2} \delta_{i, l} + l^2 h_{l-1}^{m-1} \delta_{i, l}] \delta_{n, m} \\ &\quad + [(l+1)(m-l)^2 h_l^{m-1} \delta_{i-1, l} + l^2(m+1-l) h_{l-1}^{m-1} \delta_{i, l}] \delta_{n, m+1}. \end{aligned}$$

In particular,

$$\begin{aligned} \langle Q_0^n, Q_l^m \rangle_\nabla &= (m-1)^2 h_0^{m-2} \delta_{l, 1} \delta_{n, m-1} + m^2 h_0^{m-1} \delta_{l, 0} \delta_{n, m}, \\ \langle Q_n^n, Q_l^m \rangle_\nabla &= (m-1)^2 h_{m-2}^{m-2} \delta_{l, n} \delta_{n, m-1} + m^2 h_{m-1}^{m-1} \delta_{l, n} \delta_{n, m}. \end{aligned}$$

Proof. Directly from the definition,

$$\langle Q_i^n, Q_l^m \rangle_{\nabla} = \langle \nabla Q_i^n, \nabla Q_l^m \rangle = \langle \partial_1 Q_i^n, \partial_1 Q_l^m \rangle + \langle \partial_2 Q_i^n, \partial_2 Q_l^m \rangle.$$

By Lemmas 2.4 and 3.1, the inner product $\langle \partial_j Q_i^n, \partial_j Q_l^m \rangle$ can be computed by the orthogonality of P_k^n and (3.1). For example,

$$\begin{aligned} \langle \partial_1 Q_i^n, \partial_1 Q_l^m \rangle &= (n-i)(m-l) \langle P_i^{n-1}, P_l^{m-1} \rangle + l(n-i)(m-l) \langle P_i^{n-1}, P_{l-1}^{m-2} \rangle \\ &\quad + i(n-i)(m-l) \langle P_{i-1}^{n-2}, P_l^{m-1} \rangle + il(n-i)(m-l) \langle P_{i-1}^{n-2}, P_{l-1}^{m-2} \rangle \\ &= (n-i)(m-l) h_i^{n-1} \delta_{i,l} \delta_{n,m} + l(n-i)(m-l) h_i^{n-1} \delta_{i,l-1} \delta_{n,m-1} \\ &\quad + i(n-i)(m-l) h_{i-1}^{n-2} \delta_{i-1,l} \delta_{n-1,m} + il(n-i)(m-l) h_{i-1}^{n-2} \delta_{i,l} \delta_{n,m} \end{aligned}$$

The other terms are computed similarly. \square

Corollary 3.3. For $0 \leq i \leq n$, $0 \leq l \leq m$, and $m \leq n-1$ it holds

$$\langle Q_i^n, Q_l^m \rangle_{\nabla} = [(l+1)(m-l)^2 h_l^{m-1} \delta_{i-1,l} + l^2(m+1-l) h_{l-1}^{m-1} \delta_{i,l}] \delta_{n-1,m}.$$

In particular,

$$\langle Q_0^n, Q_l^m \rangle_{\nabla} = 0 \quad \text{and} \quad \langle Q_n^n, Q_l^m \rangle_{\nabla} = 0, \quad m < n.$$

To determine the matrix \mathbf{A}_{n-1} , we will need explicit forms of the following two matrices:

$$\mathbf{C}_n := \langle \mathbb{Q}_{n+1}, \mathbb{Q}_n^{\mathsf{T}} \rangle_{\nabla} \quad \text{and} \quad \mathbf{D}_n := \langle \mathbb{Q}_n, \mathbb{Q}_n^{\mathsf{T}} \rangle_{\nabla}.$$

Lemma 3.4. For $n = 0, 1, 2, \dots$, \mathbf{D}_n is a diagonal matrix

$$(3.3) \quad \mathbf{D}_n = \text{diag}\{d_0^n, d_1^n, \dots, d_n^n\},$$

where

$$d_j^n = (n-j)^2 h_j^{n-1} + j^2 h_{j-1}^{n-1} + 2j^2(n-j)^2 h_{j-1}^{n-2}, \quad 0 \leq j \leq n,$$

with h_j^m as given in (3.1), and $\mathbf{C}_n : (n+2) \times (n+1)$ is a bidiagonal matrix,

$$(3.4) \quad \mathbf{C}_n = \begin{bmatrix} 0 & 0 & & \dots & 0 \\ c_{1,0}^n & c_{1,1}^n & & & \\ & c_{2,1}^n & c_{2,2}^n & & \\ & & \ddots & \ddots & \\ & & & c_{n,n-1}^n & c_{n,n}^n \\ 0 & \dots & & 0 & 0 \end{bmatrix},$$

where

$$\begin{aligned} c_{i,i}^n &= i^2(n-i+1) h_{i-1}^{n-1}, & 1 \leq i \leq n, \\ c_{i+1,i}^n &= (i+1)(n-i)^2 h_i^{n-1}, & 0 \leq i \leq n-1. \end{aligned}$$

Proof. The formula for \mathbf{D}_n follows directly from Lemma 3.2. Furthermore, by Corollary 3.3, for $1 \leq i \leq n-1$,

$$\langle \nabla Q_i^n, \nabla Q_i^{n-1} \rangle = (l+1)(n-1-l)^2 h_l^{n-2} \delta_{i,l+1} + l^2(n-l) h_{l-1}^{n-2} \delta_{i,l},$$

which shows that \mathbf{C}_n is a bidiagonal matrix and its first and the last row are zero. \square

We are now ready to determine the matrix \mathbf{A}_{n-1} in (3.2).

Theorem 3.5. Let $\mathbf{H}_n^\nabla := \langle \mathbb{S}_n, \mathbb{S}_n^\top \rangle_\nabla$. Then \mathbf{H}_n^∇ satisfies the recursive relation

$$(3.5) \quad \mathbf{H}_n^\nabla = \mathbf{D}_n - \mathbf{C}_{n-1}(\mathbf{H}_{n-1}^\nabla)^{-1}\mathbf{C}_{n-1}^\top,$$

where the iteration is initiated by $\mathbf{H}_1^\nabla = \mathbf{I}$, the identity matrix. Furthermore, for $n = 1, 2, \dots$, the matrix \mathbf{A}_n in (3.2) is determined by

$$(3.6) \quad \mathbf{A}_n = \mathbf{C}_n(\mathbf{H}_n^\nabla)^{-1}.$$

Proof. Using the orthogonality of \mathbb{S}_n and the fact that $S_k^n - Q_k^n \in \Pi_{n-1}^2$, we obtain from (3.2) that

$$\begin{aligned} \langle \mathbb{S}_{n+1}, \mathbb{S}_n^\top \rangle_\nabla &= \langle \mathbb{Q}_{n+1}, \mathbb{Q}_n^\top \rangle_\nabla - \mathbf{A}_n \langle \mathbb{S}_n, \mathbb{S}_n^\top \rangle_\nabla \\ &= \langle \mathbb{Q}_{n+1}, \mathbb{Q}_n^\top \rangle_\nabla - \mathbf{A}_n \langle \mathbb{Q}_n, \mathbb{S}_n^\top \rangle_\nabla \\ &= \langle \mathbb{Q}_{n+1}, \mathbb{Q}_n^\top \rangle_\nabla - \mathbf{A}_n \langle \mathbb{Q}_n, (\mathbb{Q}_n - \mathbf{A}_{n-1} \mathbb{S}_{n-1})^\top \rangle_\nabla, \end{aligned}$$

where we have used (3.2) once more. Hence, it follows that

$$\begin{aligned} \langle \mathbb{S}_{n+1}, \mathbb{S}_n^\top \rangle_\nabla &= \langle \mathbb{Q}_{n+1}, \mathbb{Q}_n^\top \rangle_\nabla - \mathbf{A}_n \langle \mathbb{Q}_n, \mathbb{Q}_n^\top \rangle_\nabla + \mathbf{A}_n \langle \mathbb{Q}_n, \mathbb{S}_{n-1}^\top \rangle_\nabla \mathbf{A}_{n-1}^\top \\ &= \langle \mathbb{Q}_{n+1}, \mathbb{Q}_n^\top \rangle_\nabla - \mathbf{A}_n \langle \mathbb{Q}_n, \mathbb{Q}_n^\top \rangle_\nabla + \mathbf{A}_n \langle \mathbb{Q}_n, \mathbb{Q}_{n-1}^\top \rangle_\nabla \mathbf{A}_{n-1}^\top. \end{aligned}$$

Consequently, from $\langle \nabla \mathbb{S}_{n+1}, \nabla \mathbb{S}_n^\top \rangle = 0$ we obtain

$$(3.7) \quad \mathbf{A}_n [\langle \mathbb{Q}_n, \mathbb{Q}_n^\top \rangle_\nabla - \langle \mathbb{Q}_n, \mathbb{Q}_{n-1}^\top \rangle_\nabla \mathbf{A}_{n-1}^\top] = \langle \mathbb{Q}_{n+1}, \mathbb{Q}_n^\top \rangle_\nabla.$$

Next we compute $\mathbf{H}_n^\nabla = \langle \mathbb{S}_n, \mathbb{S}_n^\top \rangle_\nabla$ by using (3.2) and the orthogonality of \mathbb{S}_n ,

$$(3.8) \quad \begin{aligned} \mathbf{H}_n^\nabla &= \langle \mathbb{Q}_n, \mathbb{S}_n^\top \rangle_\nabla = \langle \mathbb{Q}_n, (\mathbb{Q}_n - \mathbf{A}_{n-1} \mathbb{S}_{n-1})^\top \rangle_\nabla \\ &= \langle \mathbb{Q}_n, \mathbb{Q}_n^\top \rangle_\nabla - \langle \mathbb{Q}_n, \mathbb{S}_{n-1}^\top \rangle_\nabla \mathbf{A}_{n-1}^\top \\ &= \langle \mathbb{Q}_n, \mathbb{Q}_n^\top \rangle_\nabla - \langle \mathbb{Q}_n, \mathbb{Q}_{n-1}^\top \rangle_\nabla \mathbf{A}_{n-1}^\top. \end{aligned}$$

Since \mathbf{H}_n^∇ is nonsingular, substituting the above relation into (3.7) proves (3.6). Furthermore, substituting (3.6) into (3.8) shows that \mathbf{H}_n^∇ satisfies

$$\mathbf{H}_n^\nabla = \langle \mathbb{Q}_n, \mathbb{Q}_n^\top \rangle_\nabla - \langle \mathbb{Q}_n, \mathbb{Q}_{n-1}^\top \rangle_\nabla (\langle \mathbb{Q}_n, \mathbb{Q}_{n-1}^\top \rangle_\nabla (\mathbf{H}_{n-1}^\nabla)^{-1})^\top,$$

which simplifies to (3.5) from the symmetry of \mathbf{H}_{n-1}^∇ , and therefore completes the proof. \square

The theorem shows that \mathbf{H}_n^∇ , hence \mathbf{A}_n , can be determined iteratively.

Since $S_0^n = Q_0^n$ and $S_n^n = Q_n^n$, we only need to determine S_k^n for $1 \leq k \leq n-1$. This additional information is reflected in the matrix structure, as shown in Theorem 2.5 and (3.4),

$$\mathbf{A}_{n-1} = \begin{bmatrix} 0 & \dots & 0 \\ \hline & \tilde{\mathbf{A}}_{n-1} & \\ \hline 0 & \dots & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{C}_{n-1} = \begin{bmatrix} 0 & \dots & 0 \\ \hline & \tilde{\mathbf{C}}_{n-1} & \\ \hline 0 & \dots & 0 \end{bmatrix},$$

where $\tilde{\mathbf{A}}_{n-1}$ and $\tilde{\mathbf{C}}_{n-1}$ are matrices of size $(n-1) \times n$. These suggest a further simplification in the iteration, which we now explore.

The matrix structure shows that

$$\mathbf{H}_n^\nabla = \mathbf{D}_n - \mathbf{C}_{n-1} \mathbf{A}_{n-1}^\top = \begin{bmatrix} d_0^n & & \\ & \tilde{\mathbf{D}}_n & \\ & & d_n^n \end{bmatrix} - \begin{bmatrix} 0 & \cdots & 0 \\ \vdots & \tilde{\mathbf{C}}_{n-1} \tilde{\mathbf{A}}_{n-1}^\top & \vdots \\ 0 & \cdots & 0 \end{bmatrix},$$

which shows that the matrix \mathbf{H}_n^∇ takes the form

$$(3.9) \quad \mathbf{H}_n^\nabla = \begin{bmatrix} d_0^n & & 0 \\ & \hat{\mathbf{H}}_n^\nabla & \\ 0 & & d_n^n \end{bmatrix} \quad \text{with} \quad \hat{\mathbf{H}}_n^\nabla = \tilde{\mathbf{D}}_n - \tilde{\mathbf{C}}_{n-1} \tilde{\mathbf{A}}_{n-1}^\top.$$

Consequently, we only need to determine $\hat{\mathbf{H}}_n^\nabla$. Let us further write

$$\tilde{\mathbf{C}}_n = \left[\begin{array}{c|c|c} c_{1,0}^n & & 0 \\ \vdots & \hat{\mathbf{C}}_n & \vdots \\ 0 & & c_{n,n}^n \end{array} \right] \quad \text{with} \quad \hat{\mathbf{C}}_n = \begin{bmatrix} c_{1,1}^n & & & & \circ \\ c_{2,1}^n & c_{2,2}^n & & & \\ & \ddots & \ddots & & \\ & & & c_{n-1,n-2}^n & c_{n-1,n-1}^n \\ \circ & & & & c_{n,n-1}^n \end{bmatrix}.$$

It then follows from $\mathbf{A}_n = \mathbf{C}_n (\mathbf{H}_n^\nabla)^{-1}$ at (3.6) that

$$\tilde{\mathbf{A}}_n = \tilde{\mathbf{C}}_n \begin{bmatrix} (d_0^n)^{-1} & \cdots & 0 \\ & (\hat{\mathbf{H}}_n^\nabla)^{-1} & \\ 0 & \cdots & (d_n^n)^{-1} \end{bmatrix} = \begin{bmatrix} 1 \\ \vdots \\ 0 \end{bmatrix} \hat{\mathbf{C}}_n (\hat{\mathbf{H}}_n^\nabla)^{-1} \begin{bmatrix} 0 \\ \vdots \\ 1 \end{bmatrix},$$

where we have used the fact that $c_{1,0}^n = d_0^n = n^2 h_0^{n-1}$ and $c_{n,n}^n = d_n^n = n^2 h_{n-1}^{n-1}$, which follow directly from their explicit formulas. Consequently, we see that $\tilde{\mathbf{A}}_n$ is of the form

$$(3.10) \quad \tilde{\mathbf{A}}_n = [\mathbf{e}_1 | \hat{\mathbf{A}}_n | \mathbf{e}_n] \quad \text{with} \quad \hat{\mathbf{A}}_n = \hat{\mathbf{C}}_n (\hat{\mathbf{H}}_n^\nabla)^{-1},$$

where $\mathbf{e}_1, \mathbf{e}_n$ are, respectively, the first and the last vector in the canonical basis of \mathbb{R}^n . Consequently, it follows that

$$\tilde{\mathbf{C}}_{n-1} \tilde{\mathbf{A}}_{n-1}^\top = d_0^{n-1} \mathbf{e}_1 \mathbf{e}_1^\top + \hat{\mathbf{C}}_{n-1} \hat{\mathbf{A}}_{n-1}^\top + d_{n-1}^{n-1} \mathbf{e}_{n-1} \mathbf{e}_{n-1}^\top.$$

We finally conclude by (3.9) that the matrix $\hat{\mathbf{H}}_n^\nabla$ satisfies the relation

$$\hat{\mathbf{H}}_n^\nabla = \hat{\mathbf{D}}_n - \hat{\mathbf{C}}_{n-1} \hat{\mathbf{A}}_{n-1}^\top,$$

where $\hat{\mathbf{D}}_n$ is the diagonal matrix

$$\hat{\mathbf{D}}_n = \tilde{\mathbf{D}}_n - d_0^{n-1} \mathbf{e}_1 \mathbf{e}_1^\top - d_{n-1}^{n-1} \mathbf{e}_{n-1} \mathbf{e}_{n-1}^\top.$$

Summing up, we have proved the following proposition.

Proposition 3.6. *Let $\hat{\mathbf{Q}}_n := (Q_1^n, \dots, Q_{n-1}^n)$ and $\hat{\mathbf{S}}_n := (S_1^n, \dots, S_{n-1}^n)$. Then $\hat{\mathbf{H}}_n^\nabla = \langle \hat{\mathbf{S}}_n, \hat{\mathbf{S}}_n^\top \rangle_\nabla$. Furthermore, for $n = 2, 3, \dots$,*

$$(3.11) \quad \hat{\mathbf{Q}}_n \stackrel{c}{=} \hat{\mathbf{S}}_n + [\mathbf{e}_1 | \hat{\mathbf{A}}_{n-1} | \mathbf{e}_{n-1}] \mathbf{S}_{n-1},$$

where the matrices $\hat{\mathbf{A}}_n$ of size $n \times (n-1)$ and $\hat{\mathbf{H}}_n^\nabla$ of size $(n-1) \times (n-1)$ are determined iteratively by

$$\hat{\mathbf{A}}_n = \hat{\mathbf{C}}_n (\hat{\mathbf{H}}_n^\nabla)^{-1} \quad \text{and} \quad \hat{\mathbf{H}}_n^\nabla = \hat{\mathbf{D}}_n - \hat{\mathbf{C}}_{n-1} \hat{\mathbf{A}}_{n-1}^\top$$

for $n = 3, 4, \dots$, with the starting point $\widehat{\mathbf{A}}_1 = 0$.

Example 3.7. In the case of $\alpha = \beta = 0$, the iterative algorithm gives

$$\begin{aligned}\widehat{\mathbf{A}}_2 &= \begin{bmatrix} 1 \\ 1 \end{bmatrix}, & \widehat{\mathbf{H}}_2 &= [2], \\ \widehat{\mathbf{A}}_3 &= \frac{1}{4} \begin{bmatrix} 5 & 1 \\ 5 & 5 \\ 1 & 5 \end{bmatrix}, & \widehat{\mathbf{H}}_3 &= \begin{bmatrix} 10 & -2 \\ -2 & 10 \end{bmatrix}, \\ \widehat{\mathbf{A}}_4 &= \frac{1}{56} \begin{bmatrix} 90 & 24 & 6 \\ 53 & 72 & 11 \\ 11 & 72 & 53 \\ 6 & 24 & 90 \end{bmatrix}, & \widehat{\mathbf{H}}_4 &= \begin{bmatrix} 93 & -12 & -3 \\ -12 & 48 & -12 \\ -3 & -12 & 93 \end{bmatrix}.\end{aligned}$$

Once the matrices $\widehat{\mathbf{A}}_n$ are determined, the relation (3.11) can be used to determine the Sobolev orthogonal polynomials \mathbb{S}_n iteratively, since

$$\widehat{\mathbb{S}}_n \stackrel{c}{=} \widehat{\mathbb{Q}}_n - Q_0^{n-1} \mathbf{e}_1 - Q_{n-1}^{n-1} \mathbf{e}_{n-1} - \widehat{\mathbf{A}}_{n-1} \widehat{\mathbb{S}}_{n-1},$$

where we have used $S_0^{n-1} = Q_0^{n-1}$ and $S_{n-1}^{n-1} = Q_{n-1}^{n-1}$.

We could also determine the polynomials S_k^n directly by solving a linear system of equations. For this purpose, we fix k , $1 \leq k \leq n-1$, write

$$(3.12) \quad S_k^n(x, y) \stackrel{c}{=} Q_k^n(x, y) + \sum_{j=1}^{n-1} \sum_{i=0}^j a_i^j Q_i^j(x, y)$$

and determine the coefficient a_i^j by the orthogonality $\langle S_k^n, Q_j^m \rangle_{\nabla} = 0$ for $0 \leq l \leq m \leq n-1$, which is equivalent to the linear system of equations

$$\sum_{j=1}^{n-1} \sum_{i=0}^j a_i^j \langle Q_i^j, Q_l^m \rangle_{\nabla} = -\langle Q_k^n, Q_l^m \rangle_{\nabla}, \quad 0 \leq l \leq m \leq n-1.$$

By Lemma 3.2, these equations become

$$\begin{aligned}l(m-l)^2 h_{l-1}^{m-2} a_{l-1}^{m-1} + l^2(m-l) h_{l-1}^{m-2} a_l^{m-1} \\ + [(m-l)^2 h_l^{m-1} + 2(m-l)l^2 h_{l-1}^{m-2} + l^2 h_{l-1}^{m-1}] a_l^m \\ + l^2(m-l+1) h_{l-1}^{m-1} a_l^{m+1} + (l+1)(m-l)^2 h_l^{m-1} a_{l+1}^{m+1} \\ = -[(l+1)(m-l)^2 h_l^{m-1} \delta_{k-1,l} + l^2(m+1-l) h_{l-1}^{m-1} \delta_{k,l}] \delta_{m,n-1}.\end{aligned}$$

Observe that for $m = n-1$ the third term in the left hand side does not appear since $a_l^n = 0$ by definition. Using $h_{l-1}^{m-1} = (m-l)(\alpha + m-l) h_{l-1}^{m-2}$ and $h_l^{m-1} = l(\beta + l) h_{l-1}^{m-2}$, the above equations can be simplified to

$$(3.13) \quad \begin{aligned}(m-l) a_{l-1}^{m-1} + l a_l^{m-1} + [l\alpha + (m-l)\beta + 4l(m-l)] a_l^m \\ + l(m-l+1)(\alpha + m-l) a_l^{m+1} + (l+1)(m-l)(\beta + l) a_{l+1}^{m+1} \\ = -[(l+1)(m-l)(\beta + l) \delta_{k,l+1} + l(m+1-l)(\alpha + m-l) \delta_{k,l}] \delta_{m,n-1}.\end{aligned}$$

The indexes of a_l^m are lattices in $\Lambda_n := \{(l, m) : 0 \leq l \leq m \leq n-1\}$. For each (l, m) , the equation (3.13) involves a_l^m and its four neighbors, directly above and

below, left and right of a_l^m in the lattice. In particular, for $l = 0$ and $l = m$ we obtain the equations

$$a_0^m + a_1^{m+1} = -\delta_{k,1} \delta_{m,n-1}, \quad a_m^m + a_m^{m+1} = -\delta_{k,m} \delta_{m,n-1}.$$

By $a_l^n = 0$, these equations can be written in an equivalent way as

$$(3.14) \quad \begin{aligned} a_0^{n-1} &= -\delta_{k,1} & a_{n-1}^{n-1} &= -\delta_{k,n-1} & a_m^m + a_m^{m+1} &= 0 \\ a_0^m + a_1^{m+1} &= 0, & 1 &\leq m \leq n-2. \end{aligned}$$

These provide the boundary relations for the lattice Λ_n . Together, (3.13) and (3.14) form a linear system of equations that can be solved for $\{a_l^m : 0 \leq l \leq m \leq n-1\}$. Furthermore, the relations in (3.14) allow us to combine some of the terms in the sum (3.12). We summarize the above consideration into the following proposition.

Proposition 3.8. *For $1 \leq k \leq n-1$, the monic Sobolev polynomials are given by*

$$\begin{aligned} S_k^n &\stackrel{c}{=} Q_k^n - \delta_{k,1} Q_0^{n-1} - \delta_{n,n-1} Q_{n-1}^{n-1} + \sum_{j=1}^{n-2} a_0^j (Q_0^j - Q_1^{j+1}) \\ &\quad + \sum_{j=1}^{n-2} a_j^j (Q_j^j - Q_j^{j+1}) + \sum_{j=4}^{n-1} \sum_{i=4}^j a_{i-2}^j Q_{i-2}^j \end{aligned}$$

where the coefficients a_i^j are solutions of (3.13) and (3.14).

Example 3.9. *For the case of $\alpha = \beta = 0$, the monic Laguerre–Sobolev orthogonal polynomials satisfy the relation*

$$S_{n-k}^n(x, y) = S_k^n(y, x), \quad 0 \leq k \leq n.$$

The following are these polynomials in lower degrees: $S_0^1(x, y) = x$,

$$\begin{aligned} S_0^2(x, y) &= x(x-2), & S_1^2(x, y) &= xy - x - y, \\ S_0^3(x, y) &= x(x^2 - 6x + 6), & S_1^3(x, y) &= x^2y - x^2 - 3xy + 3x + y. \end{aligned}$$

Remark 3.1. In the case of $\alpha = \beta = 0$ we have (see equation (5.2.1) in [8])

$$q_n(u_0; x) = (-1)^n \frac{1}{n!} L_n^{-1}(x) = (-1)^{n-1} \frac{1}{(n-1)!} x L_{n-1}^1(x),$$

and therefore the constant term in $q_n(u_0; x)$ always vanishes for $n \geq 1$. Consequently, in this case, equations that hold under *modulo constant*, or $\stackrel{c}{=}$, in Theorem 2.5 can be replaced by the usual equal sign.

4. THE PRODUCT GEGENBAUER WEIGHT

In this section we study the product of Gegenbauer (or ultraspherical) weight functions and the inner product (2.3). Let

$$u_\alpha(x) := (1-x^2)^{\alpha-\frac{1}{2}}, \quad \alpha > -\frac{1}{2}.$$

The classical Gegenbauer polynomials C_n^λ , defined by ([8, Chapt IV])

$$C_n^\alpha(x) := \binom{n+2\alpha-1}{n} {}_2F_1\left(-n, n+2\alpha; \alpha + \frac{1}{2}; x\right) = 2^n \binom{n+\alpha-1}{n} x^n + \dots$$

are orthogonal with respect to the inner product

$$\langle f, g \rangle_{u_\alpha} := \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha + 1/2)\Gamma(1/2)} \int_{-1}^1 f(x)g(x) u_\alpha(x) dx.$$

More precisely, they satisfy

$$\langle C_n^\alpha, C_m^\alpha \rangle_{u_\alpha} = \frac{2^{1-2\alpha} \alpha \sqrt{\pi}}{\Gamma(\alpha + 1/2) \Gamma(\alpha)} \frac{\Gamma(n + 2\alpha)}{(n + \alpha) n!} \delta_{n,m}.$$

Moreover, they are self-coherent since they satisfy ([8, (4.7.29) in p. 83])

$$2(n + \alpha) C_n^\alpha(x) = \frac{d}{dx} [C_{n+1}^\alpha(x) - C_{n-1}^\alpha(x)], \quad n \geq 1.$$

Monic Gegenbauer orthogonal polynomials $p_n(u_\alpha)$ are defined by

$$p_n(u_\alpha; x) := 2^{-n} \binom{n + \alpha - 1}{n}^{-1} C_n^\alpha(x),$$

and their L^2 norms are given by

$$h_n^\alpha := \langle p_n(u_\alpha), p_n(u_\alpha) \rangle_{u_\alpha} = \frac{2^{1-2\alpha-2n} \sqrt{\pi} n! \Gamma(\alpha + 1) \Gamma(n + 2\alpha)}{\Gamma(\alpha + 1/2) \Gamma(n + \alpha) \Gamma(n + \alpha + 1)}.$$

From these relations, we deduce that the polynomial

$$q_n(u_\alpha; x) := p_n(u_\alpha; x) + n b_{n-1}(\alpha) p_{n-2}(u_\alpha; x),$$

where

$$b_{n-1}(\alpha) = -\frac{(n-1)}{4(n+\alpha-1)(n+\alpha-2)}, \quad n \geq 2,$$

satisfies $q'_n(u_\alpha; x) = n p_{n-1}(u_\alpha; x)$ for $n = 1, 2, \dots$

We define the product Gegenbauer weight function $U_{\alpha,\beta}(x, y) := u_\alpha(x)u_\beta(y)$ on $[-1, 1] \times [-1, 1]$ for $\alpha, \beta > -1/2$ and define monic product polynomials

$$P_k^n(x, y) := p_{n-k}(u_\alpha; x) p_k(u_\beta; y), \quad 0 \leq k \leq n.$$

These are mutually orthogonal polynomials, and

$$(4.1) \quad h_k^n := \langle P_k^n, P_k^n \rangle_{U_{\alpha,\beta}} = h_{n-k}^\alpha h_k^\beta.$$

We also define the monic polynomial Q_k^n by

$$Q_k^n(x, y) := q_{n-k}(u_\alpha; x) q_k(u_\beta; y), \quad 0 \leq k \leq n.$$

In this setting, their partial derivatives for $1 \leq k \leq n$ in Lemma 2.4 become

Lemma 4.1. For $1 \leq k \leq n-1$,

$$\begin{aligned} \partial_1 Q_k^n(x, y) &= (n-k) [P_k^{n-1}(x, y) + k b_{k-1}(\beta) P_{k-2}^{n-3}(x, y)], \\ \partial_2 Q_k^n(x, y) &= k [P_{k-1}^{n-1}(x, y) + (n-k) b_{n-k-1}(\alpha) P_{k-1}^{n-3}(x, y)]. \end{aligned}$$

Denote by $\mathcal{V}_n^2(\nabla, U_{\alpha,\beta})$, $n \geq 1$, the space of Sobolev orthogonal polynomials with respect to the bilinear form $\langle \cdot, \cdot \rangle_\nabla$ defined in (2.4), and let $S_k^n = x^{n-k} y^k + \dots$ be the monic orthogonal polynomials in $\mathcal{V}_n^2(\nabla, U_{\alpha,\beta})$. In this case, relation (2.12) becomes

$$(4.2) \quad \mathbb{Q}_n \stackrel{c}{=} \mathbb{S}_n + \mathbf{B}_{n-2} \mathbb{S}_{n-2}.$$

To compute \mathbf{B}_{n-2} explicitly, we need explicit formulas for the inner products of the gradients of the polynomials Q_k^n . In order to simplify the expressions, from now on we will write $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_{U_{\alpha,\beta}}$.

Lemma 4.2. For $0 \leq i \leq n$ and $0 \leq l \leq m$,

$$\begin{aligned} \langle Q_i^n, Q_l^m \rangle_{\nabla} = & \delta_{n,m+2} [(m-l)^2(l+2) b_{l+1}(\beta) h_l^{m-1} \delta_{i,l+2} \\ & + l^2(m-l+2) b_{m-l+1}(\alpha) h_{l-1}^{m-1} \delta_{i,l}] \\ & + \delta_{n,m} [(m-l)^2 h_l^{m-1} \delta_{i,l} + l^2(m-l)^2 b_{l-1}^2(\beta) h_{l-2}^{m-3} \delta_{i,l} \\ & + l^2 h_{l-1}^{m-1} \delta_{i,l} + l^2(m-l)^2 b_{m-l-1}^2(\alpha) h_{l-1}^{m-3} \delta_{i,l}] \\ & + \delta_{n,m-2} [l(m-l)^2 b_{l-1}(\beta) h_{l-2}^{m-3} \delta_{i,l-2} \\ & + l^2(m-l) b_{m-l-1}(\alpha) h_{l-1}^{m-3} \delta_{i,l}]. \end{aligned}$$

In particular,

$$\begin{aligned} \langle Q_0^n, Q_l^m \rangle_{\nabla} &= 2(m-2)^2 b_1(\beta) h_0^{m-1} \delta_{l,2} \delta_{n,m-2} + m^2 h_0^{m-1} \delta_{l,0} \delta_{n,m}. \\ \langle Q_n^n, Q_l^m \rangle_{\nabla} &= 2(m-2)^2 b_1(\alpha) h_{m-1}^{m-1} \delta_{l,n} \delta_{n,m-2} + m^2 h_{m-1}^{m-1} \delta_{l,n} \delta_{n,m}. \end{aligned}$$

The proof is analogous to that of Lemma 3.2.

Corollary 4.3. For $0 \leq i \leq n$, $0 \leq l \leq m$, and $m \leq n-1$ it holds

$$\langle Q_i^n, Q_l^m \rangle_{\nabla} = \delta_{n,m+2} [(m-l)^2(l+2) b_{l+1}(\beta) h_l^{m-1} \delta_{i,l+2} + l^2(m-l+2) b_{m-l+1}(\alpha) h_{l-1}^{m-1} \delta_{i,l}].$$

In particular,

$$\langle Q_0^n, Q_l^m \rangle_{\nabla} = 0 \quad \text{and} \quad \langle Q_n^n, Q_l^m \rangle_{\nabla} = 0, \quad m < n.$$

To determine the matrix \mathbf{B}_{n-2} , we will need explicit forms of the following two matrices:

$$\mathbf{C}_n := \langle \mathbb{Q}_{n+2}, \mathbb{Q}_n^{\mathsf{T}} \rangle_{\nabla} \quad \text{and} \quad \mathbf{D}_n := \langle \mathbb{Q}_n, \mathbb{Q}_n^{\mathsf{T}} \rangle_{\nabla}.$$

Lemma 4.4. For $n = 0, 1, 2, \dots$, \mathbf{D}_n is a diagonal matrix

$$(4.3) \quad \mathbf{D}_n = \text{diag}\{d_0^n, d_1^n, \dots, d_n^n\},$$

where, for $0 \leq j \leq n$,

$$\begin{aligned} d_j^n = & (n-j)^2 h_j^{n-1} + j^2(n-j)^2 b_{j-1}^2(\beta) h_{j-2}^{n-3} \\ & + j^2 h_{j-1}^{n-1} + j^2(n-j)^2 b_{n-j-1}^2(\alpha) h_{j-1}^{n-3}, \end{aligned}$$

with h_j^m as given in (4.1), and $\mathbf{C}_n : (n+3) \times (n+1)$ is a bidiagonal matrix,

$$(4.4) \quad \mathbf{C}_n = \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 \\ 0 & c_{1,1}^n & 0 & & \\ c_{2,0}^n & 0 & c_{2,2}^n & & \\ & \ddots & \ddots & \ddots & \\ & & & 0 & c_{n,n}^n \\ 0 & \cdots & & c_{n+1,n-1}^n & 0 \\ & & & 0 & 0 \end{bmatrix},$$

where

$$\begin{aligned} c_{l,l}^n &= \langle Q_l^{n+2}, Q_l^n \rangle_{\nabla} = l^2(n-l+2) b_{n-l+1}(\alpha) h_{l-1}^{n-1}, & 0 \leq l \leq n \\ c_{l+2,l}^n &= \langle Q_{l+2}^{n+2}, Q_l^n \rangle_{\nabla} = (l+2)(n-l)^2 b_{l+1}(\beta) h_l^{n-1}, & 0 \leq l \leq n. \end{aligned}$$

Proof. The formula for \mathbf{D}_n follows directly from Lemma 4.2. Furthermore, by Corollary 4.3, for $0 \leq i \leq n+2$,

$$\langle Q_i^{n+2}, Q_l^n \rangle_{\nabla} = (l+2)(n-l)^2 b_{l+1}(\beta) h_l^{n-1} \delta_{i,l+2} + l^2(n-l+2) b_{n-l+1}(\alpha) h_{l-1}^{n-1} \delta_{i,l},$$

which shows that \mathbf{C}_n is a bidiagonal matrix and its first and last row are zero. \square

Now we can compute the matrix \mathbf{B}_{n-2} in (4.2).

Theorem 4.5. *Let $\mathbf{H}_n^{\nabla} := \langle \mathbb{S}_n, \mathbb{S}_n \rangle_{\nabla}$. Then \mathbf{H}_n^{∇} satisfies the recursive relation*

$$(4.5) \quad \mathbf{H}_n^{\nabla} = \mathbf{D}_n - \mathbf{C}_{n-2}(\mathbf{H}_{n-2}^{\nabla})^{-1} \mathbf{C}_{n-2}^{\top},$$

where the iteration is initiated by $\mathbf{H}_1^{\nabla} = \mathbf{I}$, the identity matrix, and $\mathbf{H}_2^{\nabla} = \mathbf{D}_2$. Furthermore, for $n = 1, 2, \dots$, the matrix \mathbf{B}_n in (4.2) is determined by

$$(4.6) \quad \mathbf{B}_n = \mathbf{C}_n(\mathbf{H}_n^{\nabla})^{-1}.$$

Proof. This is similar to the proof of Theorem 3.5. Using (4.2) twice we obtain

$$\begin{aligned} \langle \mathbb{S}_{n+2}, \mathbb{S}_n^{\top} \rangle_{\nabla} &= \langle \mathbb{Q}_{n+2}, \mathbb{Q}_n^{\top} \rangle_{\nabla} - \mathbf{B}_n \langle \mathbb{S}_n, \mathbb{S}_n^{\top} \rangle_{\nabla} \\ &= \langle \mathbb{Q}_{n+2}, \mathbb{Q}_n^{\top} \rangle_{\nabla} - \mathbf{B}_n \langle \mathbb{Q}_n, (\mathbb{Q}_n - \mathbf{B}_{n-2} \mathbb{S}_{n-2})^{\top} \rangle_{\nabla} \\ &= \langle \mathbb{Q}_{n+2}, \mathbb{Q}_n^{\top} \rangle_{\nabla} - \mathbf{B}_n \langle \mathbb{Q}_n, \mathbb{Q}_n^{\top} \rangle_{\nabla} + \mathbf{B}_n \langle \mathbb{Q}_n, \mathbb{Q}_{n-2}^{\top} \rangle_{\nabla} \mathbf{B}_{n-2}^{\top}. \end{aligned}$$

And from $\langle \nabla \mathbb{S}_{n+2}, \nabla \mathbb{S}_n^{\top} \rangle = 0$ we deduce

$$(4.7) \quad \langle \mathbb{Q}_{n+2}, \mathbb{Q}_n^{\top} \rangle_{\nabla} = \mathbf{B}_n [\langle \mathbb{Q}_n, \mathbb{Q}_n^{\top} \rangle_{\nabla} - \langle \mathbb{Q}_n, \mathbb{Q}_{n-2}^{\top} \rangle_{\nabla} \mathbf{B}_{n-2}^{\top}].$$

Next we compute $\mathbf{H}_n^{\nabla} = \langle \mathbb{S}_n, \mathbb{S}_n^{\top} \rangle_{\nabla}$ by using (4.2) and the orthogonality of \mathbb{S}_n ,

$$(4.8) \quad \begin{aligned} \mathbf{H}_n^{\nabla} &= \langle \mathbb{Q}_n, \mathbb{S}_n^{\top} \rangle_{\nabla} = \langle \mathbb{Q}_n, (\mathbb{Q}_n - \mathbf{B}_{n-2} \mathbb{S}_{n-2})^{\top} \rangle_{\nabla} \\ &= \langle \mathbb{Q}_n, \mathbb{Q}_n^{\top} \rangle_{\nabla} - \langle \mathbb{Q}_n, \mathbb{Q}_{n-2}^{\top} \rangle_{\nabla} \mathbf{B}_{n-2}^{\top}. \end{aligned}$$

Since \mathbf{H}_n^{∇} is nonsingular, substituting the above relation into (4.7) proves (4.6). Finally, substituting (4.6) into (4.8) shows (4.5). \square

The previous theorem shows that \mathbf{H}_n^{∇} and \mathbf{B}_n can be determined iteratively.

Since $S_0^n \stackrel{c}{=} Q_0^n$ and $S_n^n \stackrel{c}{=} Q_n^n$, we only need to determine S_k^n for $1 \leq k \leq n-1$. The matrix structure reflects this information, as shown in Theorem 2.5 and (4.4); in fact we have

$$\mathbf{B}_{n-2} = \begin{bmatrix} 0 & \dots & 0 \\ \hline & \tilde{\mathbf{B}}_{n-2} & \\ \hline 0 & \dots & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{C}_{n-2} = \begin{bmatrix} 0 & \dots & 0 \\ \hline & \tilde{\mathbf{C}}_{n-2} & \\ \hline 0 & \dots & 0 \end{bmatrix},$$

where $\tilde{\mathbf{B}}_{n-2}$ and $\tilde{\mathbf{C}}_{n-2}$ are matrices of size $(n-1) \times (n-1)$.

We now proceed as in Section 3 to simplify the iteration process.

The matrix structure reads as

$$\mathbf{H}_n^{\nabla} = \mathbf{D}_n - \mathbf{C}_{n-2} \mathbf{B}_{n-2}^{\top} = \begin{bmatrix} d_0^n & & \\ & \tilde{\mathbf{D}}_n & \\ & & d_n^n \end{bmatrix} - \begin{bmatrix} 0 & \dots & 0 \\ \vdots & \tilde{\mathbf{C}}_{n-2} \tilde{\mathbf{B}}_{n-2}^{\top} & \vdots \\ 0 & \dots & 0 \end{bmatrix},$$

which shows that the matrix \mathbf{H}_n^∇ takes the form

$$(4.9) \quad \mathbf{H}_n^\nabla = \begin{bmatrix} d_0^n & & 0 \\ & \widehat{\mathbf{H}}_n^\nabla & \\ 0 & & d_n^n \end{bmatrix} \quad \text{with} \quad \widehat{\mathbf{H}}_n^\nabla = \widetilde{\mathbf{D}}_n - \widetilde{\mathbf{C}}_{n-2} \widetilde{\mathbf{B}}_{n-2}^\top,$$

and we only need to determine $\widehat{\mathbf{H}}_n^\nabla$. If we write

$$\widetilde{\mathbf{C}}_n = \left[\begin{array}{c|c|c} 0 & & 0 \\ c_{2,0}^n & & 0 \\ \vdots & \widehat{\mathbf{C}}_n & \vdots \\ 0 & & c_{n,n}^n \\ 0 & & 0 \end{array} \right] \quad \text{with} \quad \widehat{\mathbf{C}}_n = \begin{bmatrix} c_{1,1}^n & 0 & & & \\ 0 & c_{2,2}^n & & & \\ c_{3,1}^n & 0 & & & \\ & \ddots & \ddots & \ddots & \\ & & 0 & c_{n-1,n-1}^n & \\ & & c_{n,n-2}^n & 0 & \\ & & 0 & c_{n+1,n-1}^n & \end{bmatrix},$$

then from $\mathbf{B}_n = \mathbf{C}_n (\mathbf{H}_n^\nabla)^{-1}$ at (4.6) we conclude

$$\widetilde{\mathbf{B}}_n = \widetilde{\mathbf{C}}_n \begin{bmatrix} (d_0^n)^{-1} & \dots & 0 \\ & (\widehat{\mathbf{H}}_n^\nabla)^{-1} & \\ 0 & \dots & (d_n^n)^{-1} \end{bmatrix} = \begin{bmatrix} 0 & & 0 \\ 2b_1(\beta) & & 0 \\ \vdots & \widehat{\mathbf{C}}_n (\widehat{\mathbf{H}}_n^\nabla)^{-1} & \vdots \\ 0 & & 2b_1(\alpha) \\ 0 & & 0 \end{bmatrix},$$

where we use

$$\begin{aligned} c_{2,0}^n &= 2b_1(\beta) n^2 h_0^{n-1}, & d_0^n &= n^2 h_0^{n-1}, \\ c_{n,n}^n &= 2b_1(\alpha) n^2 h_{n-1}^{n-1}, & d_n^n &= n^2 h_{n-1}^{n-1}. \end{aligned}$$

Consequently, we see that $\widetilde{\mathbf{B}}_n$ is of the form

$$(4.10) \quad \widetilde{\mathbf{B}}_n = \left[2b_1(\beta) \mathbf{e}_2 | \widehat{\mathbf{B}}_n | 2b_1(\alpha) \mathbf{e}_n \right] \quad \text{with} \quad \widehat{\mathbf{B}}_n = \widehat{\mathbf{C}}_n (\widehat{\mathbf{H}}_n^\nabla)^{-1},$$

where \mathbf{e}_2 and \mathbf{e}_n are, respectively, the second vector and the second last vector in the canonical basis of \mathbb{R}^{n+1} . Consequently, it follows that

$$\widetilde{\mathbf{C}}_{n-2} \widetilde{\mathbf{B}}_{n-2}^\top = 4b_1^2(\beta) d_0^{n-2} \mathbf{e}_2 \mathbf{e}_2^\top + \widehat{\mathbf{C}}_{n-2} \widehat{\mathbf{B}}_{n-2}^\top + 4b_1^2(\alpha) d_{n-2}^{n-2} \mathbf{e}_{n-2} \mathbf{e}_{n-2}^\top.$$

We finally conclude by (4.9) that the matrix $\widehat{\mathbf{H}}_n^\nabla$ satisfies the relation

$$\widehat{\mathbf{H}}_n^\nabla = \widehat{\mathbf{D}}_n - \widehat{\mathbf{C}}_{n-2} \widehat{\mathbf{B}}_{n-2}^\top,$$

where $\widehat{\mathbf{D}}_n$ is the diagonal matrix

$$\widehat{\mathbf{D}}_n = \widetilde{\mathbf{D}}_n - 4b_1^2(\beta) d_0^{n-2} \mathbf{e}_2 \mathbf{e}_2^\top - 4b_1^2(\alpha) d_{n-2}^{n-2} \mathbf{e}_{n-2} \mathbf{e}_{n-2}^\top.$$

Summing up, we have proved the following proposition.

Proposition 4.6. *Let $\widehat{\mathbf{Q}}_n := (Q_1^n, \dots, Q_{n-1}^n)$ and $\widehat{\mathbf{S}}_n := (S_1^n, \dots, S_{n-1}^n)$. Then $\widehat{\mathbf{H}}_n^\nabla = \langle \widehat{\mathbf{S}}_n, \widehat{\mathbf{S}}_n^\top \rangle_\nabla$. Furthermore, for $n = 3, 4, \dots$,*

$$(4.11) \quad \widehat{\mathbf{Q}}_n \stackrel{c}{=} \widehat{\mathbf{S}}_n + \left[2b_1(\beta) \mathbf{e}_2 | \widehat{\mathbf{B}}_{n-2} | 2b_1(\alpha) \mathbf{e}_{n-2} \right] \mathbb{S}_{n-2},$$

where the matrices $\widehat{\mathbf{B}}_n$ of size $n \times (n-2)$ and $\widehat{\mathbf{H}}_n^\nabla$ of size $(n-1) \times (n-1)$ are determined iteratively by

$$\widehat{\mathbf{B}}_n = \widehat{\mathbf{C}}_n (\widehat{\mathbf{H}}_n^\nabla)^{-1} \quad \text{and} \quad \widehat{\mathbf{H}}_n^\nabla = \widehat{\mathbf{D}}_n - \widehat{\mathbf{C}}_{n-2} \widehat{\mathbf{B}}_{n-2}^\top$$

for $n = 3, 4, \dots$, with the initial condition $\widehat{\mathbf{B}}_1 = 0$.

Example 4.7. In the case of $\alpha = \beta = 1$ we have $b_1(1) = -\frac{1}{8}$, and the iterative algorithm gives

$$\begin{aligned} \widehat{\mathbf{B}}_2 &= -\frac{1}{8} \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad \widehat{\mathbf{H}}_2 = \begin{bmatrix} \frac{1}{2} \end{bmatrix}, \\ \widehat{\mathbf{B}}_3 &= -\frac{1}{20} \begin{bmatrix} 1 & 0 \\ 0 & 4 \\ 4 & 0 \\ 0 & 1 \end{bmatrix}, \quad \widehat{\mathbf{H}}_3 = \frac{5}{16} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\ \widehat{\mathbf{B}}_4 &= -\frac{1}{880} \begin{bmatrix} 21 & 0 & 1 \\ 0 & 110 & 0 \\ 198 & 0 & 198 \\ 0 & 110 & 0 \\ 1 & 0 & 21 \end{bmatrix}, \quad \widehat{\mathbf{H}}_4 = \frac{1}{128} \begin{bmatrix} 21 & 0 & -1 \\ 0 & 16 & 0 \\ -1 & 0 & 21 \end{bmatrix}. \end{aligned}$$

Once the matrices $\widehat{\mathbf{B}}_n$ are determined, the relation (4.11) can be used to determine the Sobolev orthogonal polynomials $\widehat{\mathbb{S}}_n$ iteratively, since

$$\widehat{\mathbb{S}}_n \stackrel{c}{=} \widehat{\mathbb{Q}}_n - Q_0^{n-2} 2b_1(\beta) \mathbf{e}_2 - Q_{n-2}^{n-2} 2b_1(\alpha) \mathbf{e}_{n-2} - \widehat{\mathbf{B}}_{n-2} \widehat{\mathbb{S}}_{n-2},$$

where we have used $S_0^n \stackrel{c}{=} Q_0^n$ and $S_n^n \stackrel{c}{=} Q_n^n$.

Example 4.8. For the case of $\alpha = \beta = 1$, the monic Gegenbauer–Sobolev orthogonal polynomials satisfy the relation

$$S_{n-k}^n(x, y) = S_k^n(y, x), \quad 0 \leq k \leq n.$$

The following are these polynomials in lower degrees:

$$\begin{aligned} S_0^1(x, y) &= x \\ S_0^2(x, y) &= x^2, \quad S_1^2(x, y) = xy, \\ S_0^3(x, y) &= x(x^2 - \frac{3}{4}), \quad S_1^3(x, y) = (x^2 - \frac{1}{4})y, \\ S_0^4(x, y) &= x^2(x^2 - 1), \quad S_1^4(x, y) = x(x^2 - \frac{5}{8})y, \quad S_2^4(x, y) = x^2y^2 - \frac{1}{4}x^2 - \frac{1}{4}y^2. \end{aligned}$$

Remark 4.1. In contrast to the Laguerre case with $\alpha = \beta = 0$, we need the *modulo constant*, or $\stackrel{c}{=}$, in the Theorem 2.5 for the Gegenbauer case. Note, however, that this is not a real limitation, since our main goal is to construct a basis for $\mathcal{V}_n^2(S)$, for which the additive constant does not matter, as shown in Theorem 2.3.

REFERENCES

- [1] C. F. Dunkl, Y. Xu, *Orthogonal polynomials of several variables*, Encyclopedia of Mathematics and its Applications **81**, Cambridge University Press, 2001, 2nd edition, 2014.
- [2] K. H. Kwon, J. K. Lee, I. H. Jung, Sobolev Orthogonal Polynomials relative to $\lambda p(c)q(c) + \tau, p'q' >$, *Comm. Korean Math. Soc.* **12** (1997), 603–617.

- [3] H. Li, Y. Xu, Spectral approximation on the unit ball, arXiv:1310.2283.
- [4] F. Marcellán, A. Branquinho, J. C. Petronilho, Classical orthogonal polynomials: A functional approach *Acta Applicandae Mathematicae* **34** (1994) 283–303.
- [5] F. Marcellán, T. E. Pérez, M. A. Piñar, Gegenbauer-Sobolev orthogonal polynomials. In *Proceedings Conference on NonLinear Numerical Methods and Rational Approximation II*. A. Cuyt ed., Kluwer Academic Publishers. Dordrecht. 1994. 71–82.
- [6] F. Marcellán, T. E. Pérez, M. A. Piñar, Laguerre–Sobolev orthogonal polynomials, *J. Comput. Appl. Math.*, **71** (2) (1996), 245–265.
- [7] F. Marcellán, Y. Xu, On Sobolev orthogonal polynomials, arXiv:1403.6249
- [8] G. Szegő, *Orthogonal polynomials*, Amer. Math. Soc. Colloq. Publ. Vol. **23**, Amer. Math. Soc. Providence, RI, 1975. Fourth Edition.
- [9] Y. Xu, Sobolev orthogonal polynomials defined via gradient on the unit ball, *J. Approx. Theory* **152** (2008), 52–65.

(L. Fernández) DEPARTAMENTO DE MATEMÁTICA APLICADA, UNIVERSIDAD DE GRANADA, SPAIN

(F. Marcellán) INSTITUTO DE CIENCIAS MATEMÁTICAS (ICMAT) AND DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD CARLOS III DE MADRID, SPAIN

(T. E. Pérez) DEPARTAMENTO DE MATEMÁTICA APLICADA, UNIVERSIDAD DE GRANADA, SPAIN

(M. A. Piñar) DEPARTAMENTO DE MATEMÁTICA APLICADA, UNIVERSIDAD DE GRANADA, SPAIN

(Y. Xu) DEPARTMENT OF MATHEMATICS, UNIVERSTY OF OREGON, USA